

Unlocking the *in vitro* anti-inflammatory and antidiabetic potential of *Polygonum maritimum*

Maria João Rodrigues, Luísa Custódio, Andreia Lopes, Marta Oliveira, Nuno R. Neng, José M. F. Nogueira, Alice Martins, Amélia P. Rauter, João Varela & Luísa Barreira

To cite this article: Maria João Rodrigues, Luísa Custódio, Andreia Lopes, Marta Oliveira, Nuno R. Neng, José M. F. Nogueira, Alice Martins, Amélia P. Rauter, João Varela & Luísa Barreira (2017) Unlocking the *in vitro* anti-inflammatory and antidiabetic potential of *Polygonum maritimum*, *Pharmaceutical Biology*, 55:1, 1348-1357, DOI: [10.1080/13880209.2017.1301493](https://doi.org/10.1080/13880209.2017.1301493)

To link to this article: <https://doi.org/10.1080/13880209.2017.1301493>



© 2017 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 17 Mar 2017.



Submit your article to this journal [↗](#)



Article views: 952



View related articles [↗](#)



View Crossmark data [↗](#)












Citing articles: 8 View citing articles [↗](#)

RESEARCH ARTICLE



Unlocking the *in vitro* anti-inflammatory and antidiabetic potential of *Polygonum maritimum*

Maria João Rodrigues^a , Luísa Custódio^a , Andreia Lopes^a, Marta Oliveira^a , Nuno R. Neng^b ,
José M. F. Nogueira^b , Alice Martins^b , Amélia P. Rauter^b , João Varela^a  and Luísa Barreira^a 

^aCCMAR, Centro de Ciências do Mar, Universidade do Algarve, Faro, Portugal; ^bCentro de Química e Bioquímica, Departamento de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Lisboa, Portugal

ABSTRACT

Context: Several *Polygonum* species (Polygonaceae) are used in traditional medicine in Asia, Europe and Africa to treat inflammation and diabetes.

Objective: Evaluate the *in vitro* antioxidant, anti-inflammatory and antidiabetic potential of methanol and dichloromethane extracts of leaves and roots of the halophyte *Polygonum maritimum* L.

Material and methods: Antioxidant activity was determined (up to 1 mg/mL) as radical-scavenging activity (RSA) of 2,2-diphenyl-1-picrylhydrazyl (DPPH), 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid) (ABTS), copper (CCA) and iron (ICA) chelating activities and iron reducing power (FRAP). NO production was measured in lipopolysaccharide (LPS)-stimulated macrophages for 24 h at concentrations up to 100 µg/mL and antidiabetic potential was assessed by α -amylase and α -glucosidase inhibition (up to 10 mg/mL) assays. The phytochemical composition of the extracts was determined by gas chromatography-mass spectrometry (GC-MS).

Results: The methanol leaf extract had the highest activity against DPPH• (IC₅₀ = 26 µg/mL) and ABTS⁺ (IC₅₀ = 140 µg/mL), FRAP (IC₅₀ = 48 µg/mL) and CCA (IC₅₀ = 770 µg/mL). Only the dichloromethane leaf extract (LDCM) showed anti-inflammatory activity (IC₅₀ = 48 µg/mL). The methanol root (IC₅₀ = 19 µg/mL) and leaf (IC₅₀ = 29 µg/mL) extracts strongly inhibited baker's yeast α -glucosidase, but LDCM had higher rat's α -glucosidase inhibition (IC₅₀ = 2527 µg/mL) than acarbose (IC₅₀ = 4638 µg/mL). GC-MS analysis identified β -sitosterol, stigmasterol, 1-octacosanol and linolenic acid as possible molecules responsible for the observed bioactivities.

Conclusions: Our findings suggest *P. maritimum* as a source of high-value health promoting commodities for alleviating symptoms associated with oxidative and inflammatory diseases, including diabetes.

ARTICLE HISTORY

Received 1 August 2016
Revised 2 February 2017
Accepted 28 February 2017

KEYWORDS

Halophytes; antioxidant activity; oxidative stress; macrophages; nitric oxide; α -amylase; α -glucosidase



Introduction

The genus *Polygonum* (Polygonaceae) includes more than 200 species worldwide, mainly in areas of temperate climate. Several *Polygonum* species are used in traditional medicine in China and Japan to treat health disorders such as dysentery, articular pain and inflammation (Takasaki et al. 2001; Kawai et al. 2006; Fan et al. 2011). Some species are also used in traditional medicine in Europe, Africa and Asia to treat diabetes (Soumyanath 2005; Bothon et al. 2013). In Europe, approximately 36 species of *Polygonum* can be found, including *P. maritimum* L., commonly known as sea knotgrass. Sea knotgrass is a perennial halophyte herb native from the sandy coasts of Europe, Mediterranean and Black Sea, Channel Islands, England and Belgium, occurring frequently throughout the Portuguese coast (Kilinc & Karaer 1995; Caçador et al. 2013). *Polygonum maritimum* has described antioxidant and antimicrobial activities (El-Haci et al. 2013), and contains bioactive molecules such as polygonocinol, (+)-8-hydroxycalamene, octacosyl, triacontyl ferulate, arylpropane, quercetin, quercitrin, (+)-catechin, and sitosterol (Kazantzoglou et al. 2009).

Diabetes is an emerging health problem in western societies affecting more than 300 million people worldwide and is

expected to be the 7th cause of death by 2030 (Mathers & Loncar 2006; Danaei et al. 2011). Type 2 diabetes mellitus (T2DM) is mainly associated with genetics and lifestyle and encompasses more than 90% of all diabetes cases globally (Mozaffarian et al. 2009). The major characteristic of T2DM is high blood glucose level, which is caused by congenital or acquired deficiency in secretion of insulin combined with decreased responsiveness to this hormone (WHO 1999; Yarchoan & Arnold 2014). The inhibition of carbohydrate-hydrolyzing enzymes, namely α -amylase and α -glucosidase, is thus an important strategy to manage hyperglycaemia linked to T2DM by decreasing the postprandial raise in blood glucose levels (Kwon et al. 2007). Acarbose, miglitol and voglibose are clinically used compounds that target α -amylase and α -glucosidase; however, they present several side effects such as abdominal distension, flatulence and meteorism (Bischoff & Flower 1985). In this sense, there has been a growing effort to search for novel natural compounds with antidiabetic properties and reduced side effects (Kwon et al. 2007).

Hyperglycaemia found in T2DM patients may also induce metabolic disturbances leading to the development of oxidative stress and chronic inflammation states that contribute to diabetes-associated complications, namely, cardiovascular,

CONTACT Luísa Barreira  lbarreir@ualg.pt  CCMAR Centro de Ciências do Mar, Universidade do Algarve, Campus de Gambelas, 8005-139 Faro, Portugal

© 2017 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

urological, neurological, kidney and eyes disorders (American Diabetes Association 2010; Vikram et al. 2014). Oxidative stress coupled with reduced antioxidant defences enhances damage caused by free radicals, such as reactive oxygen species (ROS), and contributes to disease progression (Sabu & Kuttan 2002; Maritim et al. 2003). In this context, natural antioxidants can be useful in the prevention and/or management of oxidative stress-related disorders, including diabetes (Ruhe & McDonald 2001; Fardoun 2007). ROS also contributes to the production of pro-inflammatory cytokines and chemokines and to insulin resistance (Akash et al. 2013; Muriach et al. 2014). The role of oxidative stress and chronic inflammation in the progression of T2DM thus opens new avenues in the search for novel and combined therapies comprising the prevention of oxidative and inflammatory states (Akash et al. 2013).

As stated before, several *Polygonum* species are used in traditional medicine to treat inflammation and diabetes. However, to the best of our knowledge, there is no information regarding the anti-inflammatory and/or antidiabetic potential of the sea knotgrass. In this context, we report for the first time a comparative evaluation of the antioxidant and anti-inflammatory potential and inhibitory activity on key enzymes relevant for hyperglycaemia (α -amylase and α -glucosidase) of extracts of sea knotgrass leaves and roots. The phytochemical characterization of the extracts is also presented.

Material and methods

Chemicals, culture media and supplements

Sigma-Aldrich (Germany) supplied the 1,1-diphenyl-2-picrylhydrazyl (DPPH), 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid (ABTS) radicals, sodium nitrite, lipopolysaccharide (LPS) from *Escherichia coli*, sulfanilamide, *N*-(1-Naphthyl) ethylenediamine dihydrochloride (NED) and 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT). Folin-Ciocalteu (F-C) phenol reagent and phosphoric acid were purchased from Merck (Germany). Lonza (Belgium) provided Roswell Park Memorial Institute (RPMI) 1640 medium, fetal bovine serum (FBS), L-glutamine and penicillin/streptomycin. Additional chemicals were acquired from VWR International (Belgium).

Sample collection

Whole plants of *P. maritimum* were hand collected in Ludo, South of Portugal, in June 2013. The taxonomical classification was performed by the botanist Dr. Manuel J. Pinto from the National Museum of Natural History (University of Lisbon, Botanical Garden, Portugal) and a voucher specimen is kept in the herbarium of MarBiotech laboratory (MBH22). Plants were divided in roots and leaves, washed, oven dried for 3 days at 50 °C, powdered and stored at -20 °C.

Preparation of the extracts

Dried samples were separately extracted with methanol and dichloromethane (1:40, w/v), overnight at room temperature (RT), under stirring. The extracts were filtered (Whatman no. 4), and evaporated to dryness at 40 °C in a rotary evaporator under reduced pressure (BUCHI R-210, Flawil, Switzerland). The dried extracts were dissolved in the corresponding solvent at the concentration of 10 mg/mL to be used in the chemical characterization assays, or in dimethyl sulfoxide (DMSO) to be used in the

bioactivity assays. All samples were stored at -20 °C until needed.

Gas chromatography and mass spectrometry (GC-MS) phytochemical analysis

The extracts (100 μ L) were filtered (0.2 μ m polytetrafluoroethylene membrane syringe filters), transferred to a glass vial and the solvent evaporated under a nitrogen stream. When dried, 50 μ L of the derivatization reagent [*N*-methyl-*N*-(trimethylsilyl) trifluoroacetamide; MSTFA] was added. With the vial capped, the extracts were vortexed and heated for 20 min in a dry block heater at 40 °C (Pereira et al. 2012).

The GC/MS analyses were performed on an Agilent 6890 series gas chromatograph equipped with an Agilent 7683 automatic liquid sampler coupled to an Agilent 5973 N mass selective detector (Agilent Technologies, Little Falls, DE). A programmed temperature vaporization injector with a septumless sampling head (Gerstel, Mullheim a/d Ruhr, Germany) and a baffled liner was used, operating in the solvent vent mode with compressed air for inlet cooling. Large volume injection was performed (vent time, 0.30 min; flow, 50 mL/min; pressure, 0 psi; purge, 60 mL/min at 2 min), for which the inlet temperature was programmed from 60 °C (0.4 min) to 300 °C (3 min isothermal) at a rate of 60 °C/min and subsequently decreased to 200 °C (held until end) at a rate of 50 °C/min. The injection volume and speed were set at 10 μ L and 100 μ L/min, respectively. GC analysis was performed on a Zebron ZB-5 (30 m \times 0.25 mm I.D., 0.25 μ m df; Phenomenex, USA) capillary column (5% phenyl, 95% polydimethylsiloxane), using helium as carrier gas maintained in a constant inlet pressure mode of 7.81 psi. The oven temperature was programmed from 100 °C (1 min) at 20 °C/min to 250 °C, then at 10 °C/min to 300 °C and hold for 20 min. The transfer line, ion source and quadrupole analyzer temperatures were maintained at 280 °C, 230 °C and 150 °C, respectively and a solvent delay of 4 min was selected. In the full-scan mode, electron ionization mass spectra in the range 35–550 Da were recorded at 70 eV with an ionization current of 34.6 μ A. Data recording and instrument control were performed by MSD ChemStation software (G1701CA; version C.00.00; Agilent Technologies).

Radical-scavenging activity (RSA) on DPPH[•]

The RSA against DPPH was determined according to the method described by Custódio et al. (2015). Extracts (22 μ L at concentrations ranging from 60 to 1000 μ g/mL) were mixed with 200 μ L of DPPH solution (120 μ M, in methanol) in 96-well microplates and incubated in the dark for 30 min (RT). The absorbance was measured at 517 nm (Biotek Synergy 4) and results presented as half maximal inhibitory concentration (IC₅₀, μ g/mL). Butylated hydroxytoluene (BHT) was used as a positive control.

RSA on ABTS^{•+}

The RSA against ABTS radical was evaluated by the method described previously (Rodrigues et al. 2015). A stock solution of ABTS^{•+} (7.4 mM) was generated by reacting equal amounts of ABTS with potassium persulfate (2.6 mM) for 16 h in the dark at RT. The ABTS^{•+} solution was diluted with ethanol to obtain an absorbance of at least 0.7 at 734 nm (Biotek Synergy 4). Extracts (10 μ L at concentrations from 60 to 1000 μ g/mL) were mixed in 96-well microplates with 190 μ L of ABTS^{•+} solution. After 6 min

of incubation the absorbance was measured at 734 nm (Biotek Synergy 4). Results were expressed IC₅₀ values (µg/mL). BHT was used as positive control.

RSA on nitric oxide (NO[•])

The NO[•] scavenging activity was evaluated according to Rodrigues et al. (2015) on extracts at concentrations between 60 and 1000 µg/mL. Samples (50 µL) were mixed with 50 µL of 10 mM sodium nitroprusside in phosphate buffer (PBS) and incubated for 90 min at RT. After, 50 µL of Griess reagent (1% of sulfanilamide and 0.1% of naphthylethylenediamine in 2.5% HPO₃) were added. The absorbance was read at 546 nm, and results were expressed as IC₅₀ values (µg/mL). N^ω-Nitro-L-arginine methyl ester hydrochloride (L-NAME) was used as standard.

Copper (Cu²⁺) chelating activity (CCA)

The CCA was assessed using pyrocatechol violet as described previously (Rodrigues et al. 2015). The extracts (30 µL) were applied at concentrations from 60 to 1000 µg/mL and mixed with 200 µL of 50 mM Na acetate buffer (pH 6), 6 µL of pyrocatechol violet (4 mM) in the above buffer and 100 µL of CuSO₄·5H₂O (50 µg/mL, in water). The change in the colour of the solution was measured at 632 nm using a microplate reader (Biotek Synergy 4). Results were expressed as IC₅₀ values (µg/mL). Ethylenediamine tetraacetic acid (EDTA) was used as a positive control.

Iron (Fe²⁺) chelating activity (ICA)

The ICA chelating activity was determined by measuring the formation of the Fe²⁺ ferrozine complex (Megías et al. 2009), according to Rodrigues et al. (2015). Extracts (30 µL at concentrations between 60 and 1000 µg/mL) were mixed in 96-well microplates with 200 µL of dH₂O and 30 µL of a FeCl₂ solution (0.1 mg/mL in water). After 30 min, 12.5 µL of ferrozine solution (40 mM in water) was added. The change in colour was measured in a microplate reader (Biotek Synergy 4) at 562 nm, and results were expressed as IC₅₀ values (µg/mL). EDTA was used as standard.

Ferric reducing antioxidant power (FRAP) assay

The ability of the extracts to reduce Fe³⁺ was assayed by the method described by Megías et al. (2009). Extracts (50 µL) were tested at concentrations ranging from 60 to 1000 µg/mL and mixed with distilled water (50 µL) and 1% potassium ferricyanide (50 µL). After an incubation of 20 min at 50 °C, 50 µL of 10% trichloroacetic acid (w/v) and ferric chloride solution (0.1%, w/v) were added and absorbance was measured at 700 nm. Increased absorbance of the reaction mixture indicates increased reducing power. Antioxidant activity was calculated relatively to the positive control (BHT; 1000 µg/mL), and expressed as IC₅₀ values (µg/mL).

Cell culture and cell viability

RAW 264.7 macrophages were maintained in RPMI culture medium supplemented with 10% heat-inactivated FBS, 1% L-glutamine (2 mM), and 1% penicillin (50 U/mL)/streptomycin (50 µg/mL), and were maintained at 37 °C in humidified

atmosphere with 5% CO₂. Exponentially growing cells were plated in 96-well tissue plates at a concentration of 1 × 10⁴ cells/well and incubated for 24 h. Extracts were then applied at different concentrations (3 to 100 µg/mL) for 24 h. Control cells were treated with DMSO at the highest concentration used in test wells (0.5%), and cell viability was determined by the MTT colorimetric assay (Mosmann 1983). Briefly, 2 h prior to the end of the incubation period 20 µL of MTT (5 mg/mL in PBS) were added to each well and further incubated at 37 °C. Then, 150 µL of DMSO was added to each well in order to dissolve the formazan crystals and absorbance was measured at 590 nm (Biotek Synergy 4).

Measurement of NO production

The NO production was evaluated using RAW 264.7 macrophages as described by Rodrigues et al. (2014). Cells were plated at 2.5 × 10⁵ cells/mL in 96-well tissue plates and allowed to adhere overnight. Afterwards, they were treated with nontoxic concentrations of the extracts, i.e., those that allowed cellular viability higher than 80%, in serum- and phenol-free culture medium, containing 100 ng/mL of LPS, for 24 h (Nishishiro et al. 2005). NO production was assessed using the Griess assay (Miranda et al. 2001). A calibration curve was prepared with different concentrations of sodium nitrite (1.5–100 µM). Results were expressed as percentage (%) of NO production, relative to a control containing DMSO (0.5%, v/v), and as IC₅₀ values (µg/mL).

α-Amylase inhibitory activity

The α-amylase inhibitory activity was determined by the method described by Xiao et al. (2006). Samples (40 µL at concentrations ranging from 1000 to 10,000 µg/mL) were mixed in 96-well microplates with 40 µL of amylase solution (100 U/mL in 0.1 M sodium phosphate buffer, pH 7.0) and 40 µL of 0.1% starch solution (diluted in the previous buffer). After 10 min at 37 °C, 20 µL of 1 M hydrochloric acid (HCl) and 100 µL of iodide solution (5 mM iodine (I₂) + 5 mM potassium iodide (KI), in distilled water) were added and the absorbance was measured at 580 nm. Results were expressed as IC₅₀ values (µg/mL). Acarbose was used as the standard at concentrations between 250 and 10,000 µg/mL.

Baker's yeast α-glucosidase inhibitory activity

Microbial (*Saccharomyces cerevisiae*) α-glucosidase inhibitory activity was determined according to the method described by Custódio et al. (2015). Samples (50 µL at concentrations ranging from 20 and 1000 µg/mL) were mixed with 100 µL of enzyme solution (1.0 U/mL, in 0.1 M sodium phosphate buffer, pH 7.0), and incubated for 10 min at 25 °C. Then, 50 µL of 5 mM *p*-nitrophenyl-α-D-glucopyranoside (NGP; diluted in the previous buffer) were added and incubated more 5 min at 25 °C. The absorbance was recorded at 405 nm using a microplate reader (Biotek Synergy 4) and results were expressed as IC₅₀ values (µg/mL). Acarbose was used as positive control at concentrations from 250 to 10,000 µg/mL.

Rat's intestinal α-glucosidase inhibitory activity

Rat's intestinal acetone powder was used as a crude enzyme extract as an example of enzyme of mammalian origin

(Kwon et al. 2007). Rat's intestinal acetone powder (250 mg) were mixed with 10 mL of 0.1 M sodium phosphate buffer (pH 7.0) and centrifuged at $5000 \times g$ for 20 min at 4 °C. The supernatant (100 μ L) was mixed with the extracts (50 μ L at concentrations between 500 and 10,000 μ g/mL), and incubated for 10 min at 25 °C. Then, 50 μ L of 5 mM *p*-nitrophenyl- α -D-glucopyranoside (NGP; diluted in the previous buffer) was added and the mixture was incubated for 30 min at 37 °C. The absorbance was read at 405 nm using a microplate reader (Biotek Synergy 4), and results were expressed as IC₅₀ values (μ g/mL). Acarbose was used as positive control at concentrations from 250 to 10,000 μ g/mL.

Statistical analysis

Results were expressed as mean \pm standard error of the mean (SEM), and experiments were carried out at least in triplicate. Significant differences were assessed by analysis of variance (ANOVA) followed by Duncan's New Multiple Range Test, or by Kruskal–Wallis test when parametricity of data did not prevail. SPSS statistical package for Windows (release 15.0, SPSS Inc.) was used. The IC₅₀ values were calculated by sigmoidal fitting of the data in the GraphPad Prism v. 5.0 software.

Results

Phytochemical analysis

In order to determine their chemical composition, extracts were analyzed by GC-MS (Figure 1 and Table 1). This analysis was able to identify a large number of compounds detected in the methanol extracts (70–75%). However, a large percentage of compounds detected in the dichloromethane extracts could not be identified (45–48%). In total, 51 compounds were identified belonging to different classes of biochemicals: alkanes and alkenes (AA), fatty acids (FA), phenolic compounds (PC), acylglycerols (GLY), saccharides (SAC), alcohols (ALC), phytosterols (PS) and minor groups (MG).

With similar contents, fatty acids (24.3%) and acylglycerols (22.8%) were the most represented categories in the methanol leaf extract (LM), in which 2-monostearin (35) was the major compound (15.4%). Palmitic acid (18), linolenic acid (23), 1-monopalmitin (34), glycerol (43), and β -sitosterol (47) were also detected at abundances higher than 5%. Acylglycerols were the most abundant constituents of the LM extract (34.4%), with 1-monostearin (31) being the most representative compound (24.0%), while 1-monopalmitin (34) and oleamide (51)




Figure 1. Main chemical compound classes identified by GC/MS in the dichloromethane and methanol extracts of roots and leaves of *P. maritimum*. (A) Methanol leaf extract; (B) Methanol root extract; (C) Dichloromethane leaf extract; and (D) Dichloromethane root extract. AA: alkanes and alkenes; FA: fatty acids; PC: phenolic compounds; GLY: acylglycerols; SAC: saccharides; ALC: alcohols; PS: phytosterols; MG: minor groups; NI: non-identified compounds.

Table 1. Phytochemical composition of the methanol and dichloromethane extracts of *P. maritimum* leaves and roots, determined by GC/MS analysis.

ID	RT (min)	Compound	Molecular formula	Relative abundance (%)			
				Methanol		Dichloromethane	
				Leaves	Roots	Leaves	Roots
<i>Alkanes/Alkenes</i>							
1	5.61	2,6,11-Trimethyl dodecane	C ₁₅ H ₃₂		0.67		
2	5.63	Phytane	C ₂₀ H ₄₂			0.22	0.21
3	5.66	2,6,10,15-Tetramethyl heptadecane	C ₁₇ H ₃₆		0.72		
4	6.75	Tetradecane	C ₁₄ H ₃₀		0.97	0.32	0.40
5	8.55	Cetane	C ₁₆ H ₃₄			0.16	0.18
6	8.63	2-Methyl octadecane	C ₁₉ H ₄₀		0.78	0.26	
7	9.78	Octacosane	C ₂₈ H ₅₈		0.77	0.37	0.70
8	9.90	3,11-Dimethyl nonacosane	C ₃₁ H ₆₄		1.10		
9	11.57	Eicosane	C ₂₀ H ₄₂		0.90		
10	12.39	Docosane	C ₂₂ H ₄₆		1.78		1.01
11	12.40	Tricosane	C ₂₃ H ₄₈			0.73	
12	14.10	Pentacosane	C ₂₅ H ₅₂			0.59	
13	16.89	1-Nonadecene	C ₁₉ H ₃₈			0.87	
14	17.75	Nonacosane	C ₂₉ H ₆₀			1.80	
<i>Fatty acids</i>							
15	5.75	Pelargonic acid	C ₁₂ H ₁₈ O ₂	0.28			0.17
16	7.72	Lauric acid	C ₁₂ H ₂₄ O ₂	1.51		0.25	0.23
17	8.92	Myristic acid	C ₁₄ H ₂₈ O ₂	1.22		1.72	0.65
18	10.28	Palmitic acid	C ₁₆ H ₃₂ O ₂	6.40	1.46	4.49	6.53
19	11.08	Linoleic acid	C ₁₈ H ₃₂ O ₂	1.79	2.32	0.35	
20	11.18	Margaric acid	C ₁₇ H ₃₄ O ₂			0.14	0.20
21	11.75	Oleic acid	C ₁₈ H ₃₄ O ₂	4.13		2.32	3.21
22	11.85	Linolenic acid	C ₁₈ H ₃₀ O ₂			0.95	
23	11.94	Stearic acid	C ₁₈ H ₃₆ O ₂	6.71	4.27	1.83	1.38
24	13.59	Arachidic acid	C ₂₀ H ₄₀ O ₂	1.71		1.05	
25	15.32	Behenic acid	C ₂₂ H ₄₄ O ₂			1.35	2.64
26	17.10	Lignoceric acid	C ₂₄ H ₄₈ O ₂	0.53		1.29	1.78
<i>Phenols</i>							
27	5.09	Benzoic acid	C ₇ H ₇ O ₂	3.24	0.51		
28	6.94	Butylated hydroxytoluene	C ₁₅ H ₂₄ O	0.99	1.31	0.17	0.27
29	7.12	Vanillin	C ₈ H ₈ O ₃				0.68
30	7.70	Phloroglucinol	C ₆ H ₆ O ₃		0.92		
31	9.63	Methyl 3-(3,4-di-tert-butyl-4-hydroxyphenyl)propionate	C ₁₈ H ₂₈ O ₃	0.68	0.66		
32	9.69	Gallic acid	C ₇ H ₆ O ₅	1.15		0.54	1.97
<i>Acylglycerols</i>							
33	14.63	2-Monopalmitin	C ₁₉ H ₃₈ O ₄				1.28
34	14.89	1-Monopalmitin	C ₁₉ H ₃₈ O ₄	5.89	10.3	2.20	3.05
35	16.14	2-Monostearin	C ₁₉ H ₃₈ O ₄	15.4		0.34	
36	16.27	1-Monoolein	C ₂₁ H ₄₀ O ₄	0.35		0.47	
37	16.59	1-Monostearin	C ₁₉ H ₃₈ O ₄	1.08	24.1	2.87	5.44
<i>Saccharides</i>							
38	8.69	D-Fructose	C ₆ H ₇ O ₆	4.40	0.97	0.39	0.27
39	9.02	β-D-Glucofuranose	C ₆ H ₁₂ O ₆	0.85			
40	9.34	β-D-Glucopyranose	C ₆ H ₁₂ O ₆	1.05		0.14	
41	9.41	α-D-Glucopyranose	C ₆ H ₁₂ O ₆	0.45			
<i>Alcohols</i>							
42	4.12	Butane-1,3-diol	C ₄ H ₁₀ O ₂	0.62			
43	5.18	Glycerol	C ₃ H ₈ O ₃	5.20	5.02		
44	11.35	Phytol	C ₂₀ H ₄₀ O	0.62			
45	20.91	1-Octacosanol	C ₂₈ H ₅₈ O			8.72	
<i>Phytosterols</i>							
46	24.51	Stigmasterol	C ₂₉ H ₄₈ O			1.37	
47	25.88	β-Sitosterol	C ₂₉ H ₅₀ O	5.90	1.88	10.6	19.0
<i>Minor groups</i>							
48	5.25	Phosphoric acid	H ₃ PO ₄	0.62	0.55	0.46	0.34
49	7.44	Diethyl phthalate	C ₁₂ H ₁₄ O ₄	0.35			
50	7.78	Thymine	C ₅ H ₆ N ₂ O ₂		0.46		
51	13.11	Oleamide	C ₁₈ H ₃₅ NO	0.89	6.66	2.30	3.13

represented 10.3 and 6.66% of the total identified components, respectively. In the dichloromethane leaf extract (LDCM), fatty acids represented 15.7% and phytosterols (PS) 12.0% of the total identified compounds. β-Sitosterol (47; 10.5%) and 1-octacosanol (45; 8.7%) were the major components identified. Similarly, the dichloromethane root extract (RDCM) had the highest content in phytosterols (PS; 19.0%) and fatty acids (FA; 16.7%), in which β-sitosterol (47, 19.0%) was the main component, followed by

palmitic acid (18) and 1-monostearin (37) at 6.5 and 5.4%, respectively.

Antioxidant activity

The *P. maritimum* LM and methanol root (RM) extracts showed high RSA for both DPPH and ABTS radicals, coupled with a

Table 2. Radical scavenging activity (RSA) on DPPH, ABTS and NO radicals, metal chelating activity on iron (ICA) and copper (CCA) and ferric reducing activity (FRAP) activity of methanol (MeOH) and dichloromethane (DCM) extracts of roots and leaves of *P. maritimum*.

Extract/Standard	Plant organ	RSA			Metal chelation/reduction		
		DPPH	ABTS	NO	CCA	ICA	FRAP
MeOH	Leaves	26 ± 0.7 ^a	140 ± 6 ^a	n.a.	290 ± 10 ^b	n.a.	48 ± 1.4 ^a
	Roots	27 ± 0.0 ^a	192 ± 5 ^b	n.a.	446 ± 8 ^c	n.a.	64 ± 3.8 ^a
DCM	Leaves	n.a.	n.a.	n.a.	n.a.	n.a.	798 ± 36 ^b
	Roots	n.a.	n.a.	n.a.	n.a.	n.a.	770 ± 53 ^b
BHT*	–	110 ± 10 ^b	140 ± 10 ^a	–	–	–	–
L-NAME*	–	–	–	2500 ± 10	–	–	–
EDTA*	–	–	–	–	170 ± 10 ^a	55.9 ± 3.7	–

Results are expressed as IC₅₀ values (µg/mL). *Positive control; n.a.: not active. Values are means ± SEM of three separate experiments performed in triplicate (n = 9). In the same column, means labelled with different letters are significantly different by Duncan's multiple range test (*p* < 0.05).

strong capacity for reducing iron and chelating copper (Table 2). In the DPPH assay, the methanol extracts of both organs had similar IC₅₀ values (roots: 27 µg/mL; leaves: 26 µg/mL), which were significantly lower than the one obtained with the positive control (BHT, 110 µg/mL; Table 2). Regarding the ABTS radical, both the RM and LM extracts had IC₅₀ of 192 and 140 µg/mL, respectively, which were similar to that of BHT (140 µg/mL; Table 2). The RM and LM had also high ability to reduce iron with IC₅₀ values of 64 and 48 µg/mL, respectively (Table 2). A moderate iron reduction was obtained upon addition of the RDCM (IC₅₀ = 770 µg/mL) and LDCM (IC₅₀ = 798 µg/mL) extracts. However, none of the extracts had the capacity for scavenging the NO radical or for chelating iron (Table 2).

Anti-inflammatory activity

To assess the *in vitro* anti-inflammatory activity, the effect of applying nontoxic concentrations of the extracts (i.e. yielding cell viability >80%) on the NO production by LPS-stimulated RAW 264.7 macrophage cells was determined. A significant reduction in cell viability was observed upon applying the LM extract at a concentration of 100 µg/mL to the cells. Loss of viability was also observed with the RDCM extract at 50 and 100 µg/mL (data not shown). Therefore, these concentrations were not used in the anti-inflammatory activity assessment.

Exposure of RAW 264.7 cells to LPS at 100 ng/mL increased the nitrite concentration in the culture medium from a basal level of approximately 0.3 µM to around 13 µM (data not shown). This increase was significantly reduced in a dose-dependent manner by the treatment with LDCM, at concentrations ranging from 25 to 100 µg/mL, showing that the latter had an activity similar to the positive control (L-NAME) at the same concentration (Figure 2). However, this extract had an IC₅₀ value of 48 µg/mL, higher than that of L-NAME (29.1 µg/mL; data not shown). Interestingly, incubating this cell line with the remaining extracts resulted in an increase in NO production when compared to the control (Figure 2). In particular, incubation with LM and RM extracts resulted in the most significant increases in NO production: 144% at the concentration of 50 µg/mL and 139% at 25 µg/mL, respectively.

Antidiabetic activity

The inhibitory potential of extracts from *P. maritimum* was evaluated against α-amylase and baker's yeast and rat's α-glucosidases (Table 3). The methanol extracts had the highest capacity to inhibit the baker's yeast α-glucosidase with IC₅₀ values of

19 and 29 µg/mL for roots and leaves, respectively, which were significantly lower than that of acarbose (IC₅₀ = 3144 µg/mL). Despite the fact that methanol extracts had no capacity to inhibit rat's α-glucosidase, LDCM had the highest rat's α-glucosidase inhibitory activity (IC₅₀ = 2527 µg/mL), which was a significantly lower IC₅₀ value than that of acarbose (4638 µg/mL; *p* < 0.05).

Discussion

In order to confirm the use of *Polygonum* species in traditional medicine in the treatment of inflammation and diabetes, we assessed for the first time the phytochemical composition of *P. maritimum* and its *in vitro* anti-inflammatory and antidiabetic potential.

GC-MS analysis detected alkanes, alkenes and fatty acids as the major categories of compounds in *P. maritimum* extracts. These are lipophilic compounds widely distributed in plants as constituents of plant waxes and have been described in the halophilic Mediterranean *Suaeda vera* Forssk. ex J.F.Gmel. (Amaranthaceae), *Sarcocornia fruticosa* (L.) A.J.Scott (Amaranthaceae), and *Halimione portulacoides* (L.) Aellen (Amaranthaceae) (Grossi & Raphael 2003). Waxes usually have a protective role, for example, against microbial infections and avoiding excessive water losses and acute osmotic stress that halophilic species suffer while growing in drenched soils (Müller & Riederer 2005; Huang et al. 2011). Moreover, they are common constituents of essential oils of halophyte species, such as *Suaeda fruticosa* and *Limonium echioides* (L.) Mill. (Plumbaginaceae) (Saïdana et al. 2008). From the compounds detected in this study, only β-sitosterol (47) had been previously identified in *P. maritimum* also in dichloromethane extracts. Phytol (44), palmitic (18) and lauric (16) acids were also previously detected as components of essential oils of *Polygonum* species, namely *P. hydropiper* L. and *P. minus* Huds. (Miyazawa & Tamura 2007; Baharum et al. 2010).

Our findings showed that *P. maritimum* has a strong antioxidant activity comparable to that reported by other authors for similar extracts made from aerial parts of the same species (El-Haci et al. 2013) and *P. sachalinensis* F.Schmidt and *P. cuspidatum* Siebold & Zucc. (Pan et al. 2007; Fan et al. 2011). Moreover, IC₅₀ values obtained with the RM and LM were significantly lower than or similar to the one obtained with the positive control (BHT) for DPPH and ABTS assays, respectively. The significantly higher antioxidant activity of these extracts can be related to the presence of some of the compounds identified in these extracts, namely, the phenolic compounds benzoic acid, BHT, vanillin and phloroglucinol, since these compounds are




Figure 2. Effect of the application of dichloromethane and methanol extracts of roots and leaves of *P. maritimum* on NO production (%) by LPS-stimulated macrophages. Control cells were treated with culture medium supplemented with 0.5% DMSO and 100 ng/mL of LPS. L-NAME (positive control) was applied at the concentration of 100 µg/mL. Solid and errors bars represent the average and SEM, respectively ($n = 9$). Bars followed by different letters are significantly different according to the Duncan's multiple ranges test ($p < 0.05$).

Table 3. Inhibitory activity of dichloromethane and methanol extracts of roots and leaves of *P. maritimum* on α -amylase, baker's yeast α -glucosidase and rat's intestinal α -glucosidase.

Extract/Standard	Organ	α -Amylase	Yeast α -glucosidase	Rat α -glucosidase
Methanol	Leaves	n.a.	29 ± 0.7 ^a	n.a.
	Roots	n.a.	19 ± 0.5 ^a	n.a.
Dichloromethane	Leaves	n.a.	585 ± 27 ^b	2527 ± 37
	Roots	n.a.	626 ± 14 ^b	>2500
Acarbose*		7797 ± 98	3144 ± 132 ^c	4638 ± 438

Results are expressed as IC₅₀ values (µg/mL). *Positive control; n.a.: not active. Values are means ± SEM of three separate experiments performed in triplicate ($n = 9$). In the same column, means labelled with different letters are significantly different by Duncan's multiple range test ($p < 0.05$).

well described as strong antioxidants *in vitro* (Foti 2007; Dai & Mumper 2010). Phytol and linolenic acid have also been described as potent *in vitro* and *in vivo* antioxidants due to their hydroxyl group (Richard et al. 2008; Santos et al. 2013). Thus, their presence in the LM could also contribute to the high RSA and ICA of this extract. Interestingly, no significant differences were observed between the antioxidant activity of leaves and roots as opposed to that reported for other halophytes species, such as *Mesembryanthemum edule* L. (Aizoaceae), *Limoniastrum monopetalum* (L.) Boiss. (Plumbaginaceae), *Salsola kali* L. (Chenopodiaceae), *Tamarix gallica* L. (Tamaricaceae) and *Limonium algarvense* Erben (Plumbaginaceae) (Ksouri et al. 2008; Falleh et al. 2012; Trabelsi et al. 2012; Rodrigues et al. 2015). This can be explained by the similar phytochemical profile of the methanol extracts of both organs (Figure 1 and Table 1), suggesting that the bioactive compound(s) are not organ-specific.

LPS is a cell wall endotoxin produced by Gram-negative bacteria that activates macrophages to produce inflammatory mediators such as NO (Martich et al. 1993), a radical associated with chronic inflammation (Kubes 2000; Joo et al. 2014). In this context, a decrease in NO production is used as an indicator of the potential for the extracts to reduce an inflammatory response (Joo et al. 2014; Rodrigues et al. 2014). On the other hand,

an increase in NO production can indicate an immunostimulatory effect of the extracts, which is important in macrophage defence and protection against infection (Wink et al. 2011).

The anti-inflammatory effect of *P. maritimum* leaves may be attributed to the presence of compound(s) with potential anti-inflammatory properties, most likely β -sitosterol, stigmasterol, 1-octacosanol, oleamide as well as linolenic and oleic acids (Table 1). For instance, β -sitosterol, one of the major constituents of this extract, has anti-inflammatory properties through TNF- α inhibition (Loizou et al. 2010), and so do 1-octacosanol and oleic acid (Vassiliou et al. 2009; de Oliveira et al. 2012). In turn, linolenic acid and oleamide are able to reduce NO production and inducible nitric oxide synthase (iNOS) gene expression, through the inhibition of the NF- κ B pathway (Ren & Chung 2007; Oh et al. 2010). Stigmasterol was also reported to inhibit the production of pro-inflammatory mediators associated with the same pathway (Gabay et al. 2010). The combination of all these compounds in the LDCM of *P. maritimum* are probably contributing to its NO inhibitory capacity, since they were solely identified in this extract, or in higher quantities (Figure 2). These findings are in accordance with several reports on the anti-inflammatory properties of *Polygonum* species (*P. lapathifolium* L., *P. cuspidatum* and *P. perfoliatum* L. (Takasaki et al. 2001; Kim et al. 2007; Fan et al. 2011; Lei et al. 2015).

NO has an important role in the immune system modulation, being one of the macrophage-mediated primary responses against pathogens such as fungi, helminthes, protozoa and bacteria (Wink et al. 2011). In this sense, the increased NO production suggests that both the LM and RM from *P. maritimum* may have immunostimulatory properties (Wink et al. 2011) most likely due to the presence of specific molecules in these extracts, namely, saturated fatty acids (stearic and palmitic acids), which are known inducers of the production of pro-inflammatory cytokines in macrophages (Valdearcos et al. 2012; Miao et al. 2015). In fact, immunostimulatory properties are reported in

P. multiflorum, *P. minus* and *P. cuspidatum* (Chen et al. 2012; Veerasamy et al. 2014; Chueh et al. 2015).

Compounds with the capacity to inhibit carbohydrate-hydrolyzing enzymes like α -amylase and α -glucosidase can delay the digestion of carbohydrates, decreasing the postprandial increase of blood glucose level after a mixed carbohydrate meal, and therefore can be an important strategy in managing hyperglycaemia linked to T2DM (Krentz & Bailey 2005; Kwon et al. 2007; Bhandari & Ansari 2008).

It is noteworthy that methanol extracts were approximately 100-fold more active towards baker's yeast α -glucosidase than acarbose and that the LDCM had 2-times more ability to inhibit rat's α -glucosidase. High IC_{50} values for acarbose against this enzyme have also been reported by other authors (4823 μ g/mL) (Gao et al. 2013). However, the methanol extracts had no capacity to inhibit rat's α -glucosidase, which is a common feature of molecules with inhibitory capacity on α -glucosidase from microbial origin (Oki et al. 1999; Shai et al. 2011). Furthermore, a few *in vitro* studies have discussed the low capacity of acarbose to inhibit mammalian α -glucosidase compared to crude extracts, including aqueous ethanol extracts from *P. senegalensis* (Meisn.) Soják (Polygonaceae), which is used in folk medicine to treat T2DM (Shinde et al. 2008; Bothon et al. 2013). Those differences are usually attributed to additive or synergistic interactions of the compounds present in the extracts, resulting in a higher capacity to inhibit the mammalian α -glucosidase (Adisakwattana et al. 2012). The higher activity of the LDCM can be related with the presence of some particular compounds. For example, β -sitosterol was the main compound identified in this extract (Table 1) and was previously reported to possess strong hypoglycaemic activity through α -glucosidase inhibition (Ortiz-Andrade et al. 2007). The same properties were described for stigmaterol and linolenic acid, present only in this extract, as well as for oleic acid (Ortiz-Andrade et al. 2007; Lean Teik et al. 2013; Su et al. 2013). In fact, Su et al. (2013) reported that linolenic and oleic acids were more active than acarbose. Although none of the extracts achieved 50% of inhibitory activity in the α -amylase assay, our data suggest that *P. maritimum* may have potential as a source of antidiabetic molecules. This is in accordance with the antidiabetic activity found in several *Polygonum* species, such as *P. aviculare* L. (Polygonaceae), *P. cuspidatum*, *P. multiflorum* and *P. senegalensis* (Soumyanath 2005; Bothon et al. 2013).

Conclusions

The present study highlights for the first time the potential of the halophyte *P. maritimum* as a source of compounds with anti-oxidant, anti-inflammatory and antidiabetic activities. The methanol extracts had the highest antioxidant capacity, possibly due to the presence of benzoic acid, BHT, phloroglucinol, phytol and linolenic acid. The dichloromethane extracts from *P. maritimum* leaves had significant anti-inflammatory activity, most likely related to its main constituents identified as β -sitosterol, stigmaterol, 1-octacosanol, oleamide, linolenic and oleic acids. Moreover, its high α -glucosidase inhibitory activity may be related to the presence of β -sitosterol, stigmaterol, linolenic and oleic acid. Overall, our results indicate that *P. maritimum* extracts are endowed with compounds with potential to be used as a combined strategy to manage T2DM due to its anti-inflammatory, antioxidant and α -glucosidase inhibitory properties. These results could be the starting points to further explore *P. maritimum*, especially its leaves, as a source of value-added bioactive natural products. Nonetheless, isolation and

identification of the molecule(s) responsible for the detected biological activities is already being pursued.

Acknowledgments

The authors thank the Faculty of Pharmacy and Center for Neurosciences and Cell Biology (University of Coimbra, Portugal) for kindly provide the murine leukemic monocyte-macrophage cell line (RAW 264.7).










Disclosure statement

The authors report that they have no conflicts of interest.

Funding

This work was supported by the XtremeBio project (PTDC/MAR-EST/4346/2012) funded by Foundation for Science and Technology (FCT) and the Portuguese National Budget. This work benefited also from national funding through FCT project CCMAR/Multi/04326/2013. Luísa Custódio was supported by the FCT Investigator Program (IF/00049/2012). Nuno R. Neng and José M. F. Nogueira also acknowledge a FCT Post-Doc grant (SFRH/BPD/86071/2012) and funding (UID/Multi/00612/2013).

ORCID

Maria João Rodrigues  <http://orcid.org/0000-0001-8732-710X>
 Luísa Custódio  <http://orcid.org/0000-0003-4338-7703>
 Marta Oliveira  <http://orcid.org/0000-0001-6793-8026>
 Nuno R. Neng  <http://orcid.org/0000-0001-9879-1565>
 José M. F. Nogueira  <http://orcid.org/0000-0001-5785-6780>
 Alice Martins  <http://orcid.org/0000-0002-0542-2662>
 Amélia P. Rauter  <http://orcid.org/0000-0003-3790-7952>
 João Varela  <http://orcid.org/0000-0003-3101-693X>
 Luísa Barreira  <http://orcid.org/0000-0002-4077-855X>

References

- Adisakwattana S, Ruengsamran T, Kampa P, Sompong W. 2012. *In vitro* inhibitory effects of plant-based foods and their combinations on intestinal α -glucosidase and pancreatic α -amylase. BMC Complement Altern Med. 12:110.
- Akash MS, Rehman K, Chen S. 2013. Role of inflammatory mechanisms in pathogenesis of type 2 diabetes mellitus. J Cell Biochem. 114:525–531.
- American Diabetes Association. 2010. Standards of Medical Care in Diabetes. Diabetes Care 33:S11–S61.
- Baharum SN, Bunawan H, Ghani MA, Mustapha WAW, Noor NM. 2010. Analysis of the chemical composition of the essential oil of *Polygonum minus* Huds. using two-dimensional gas chromatography-time-of-flight mass spectrometry (GC-TOF MS). Molecules. 15:7006–7015.
- Bhandari U, Ansari MN. 2008. Antihyperglycaemic activity of aqueous extract of *Embelia ribes* Burm in streptozotocin-induced diabetic rats. Indian J Exp Biol. 46:607–613.
- Bischoff PM, Flower RW. 1985. Ten years experience with choroidal angiography using indocyanine green dye: a new routine examination or an epilogue? Doc Ophthalmol. 60:235–291.
- Bothon FTD, Debiton E, Avlessi F, Forestier C, Teulade JC, Sohounhloue DKC. 2013. *In vitro* biological effects of two anti-diabetic medicinal plants used in Benin as folk medicine. BMC Complement Altern Med. 13:51.
- Caçador I, Neto JM, Duarte B, Barroso DV, Pinto M, Marques JC. 2013. Development of an Angiosperm Quality Assessment Index (AQuA-Index) for ecological quality evaluation of Portuguese water bodies—A multi-metric approach. Ecol Indic. 25:141–148.

- Chen Q, Zhang SZ, Ying HZ, Dai XY, Li XX, Yu CH, Ye HC. 2012. Chemical characterization and immunostimulatory effects of a polysaccharide from *Polygoni multiflori* Radix Praeparata in cyclophosphamide-induced anemic mice. *Carbohydr Polym.* 88:1476–1482.
- Chueh FS, Lin JJ, Lin JH, Weng SW, Huang YP, Chung JG. 2015. Crude extract of *Polygonum cuspidatum* stimulates immune responses in normal mice by increasing the percentage of Mac-3-positive cells and enhancing macrophage phagocytic activity and natural killer cell cytotoxicity. *Mol Med Rep.* 11:127–132.
- Custódio L, Patarra J, Alberício F, Neng NR, Nogueira JMF, Romano A. 2015. Phenolic composition, antioxidant potential and *in vitro* inhibitory activity of leaves and acorns of *Quercus suber* on key enzymes relevant for hyperglycemia and Alzheimer's disease. *Ind Crops Prod.* 64:45–51.
- Dai J, Mumper RJ. 2010. Plant phenolics: extraction, analysis and their antioxidant and anticancer properties. *Molecules.* 15:7313–7352.
- Danaei G, Finucane MM, Lu Y, Singh GM, Cowan MJ, Paciorek CJ, Lin JK, Farzadfar F, Khang YH, Stevens GA, Global burden of metabolic risk factors of chronic diseases collaborating group (blood glucose), et al. 2011. National, regional, and global trends in fasting plasma glucose and diabetes prevalence since 1980: systematic analysis of health examination surveys and epidemiological studies with 370 country-years and 2.7 million participants. *Lancet.* 378:31–40.
- de Oliveira AM, Conserva LM, de Souza Ferro JN, de Almeida Brito F, Lyra Lemos RP, Barreto E. 2012. Antinociceptive and anti-inflammatory effects of octacosanol from the leaves of *Sabicea grisea* var. *grisea* in mice. *Int J Mol Sci.* 13:1598–1611.
- El-Haci IA, Bekkara FA, Mazari W, Hassani F, Didi MA. 2013. Screening of biological activities of *Polygonum maritimum* L. from Algerian coast. *Asian Pac J Trop Biomed.* 3:611–616.
- Falleh H, Jalleli I, Ksouri R, Boulaaba M, Guyot S, Magné C, Abdelly C. 2012. Effect of salt treatment on phenolic compounds and antioxidant activity of two *Mesembryanthemum edule* provenances. *Plant Physiol Biochem.* 52:1–8.
- Fan D, Zhou X, Zhao C, Chen H, Zhao Y, Gong X. 2011. Anti-inflammatory, antiviral and quantitative study of quercetin-3-O- β -D-glucuronide in *Polygonum perfoliatum* L. *Fitoterapia.* 82:805–810.
- Fardoun RZ. 2007. The use of vitamin E in type 2 diabetes mellitus. *Clin Exp Hypertens.* 29:135–148.
- Foti MC. 2007. Antioxidant properties of phenols. *J Pharm Pharmacol.* 59:1673–1685.
- Gabay O, Sanchez C, Salvat C, Chevy F, Breton M, Nourissat G, Wolf C, Jacques C, Berenbaum F. 2010. Stigmasterol: a phytosterol with potential anti-osteoarthritic properties. *Osteoarthritis Cartilage.* 18:106–116.
- Gao J, Xu P, Wang Y, Wang Y, Hochstetter D. 2013. Combined effects of green tea extracts, green tea polyphenols or epigallocatechin gallate with acarbose on inhibition against α -amylase and α -glucosidase *in vitro*. *Molecules.* 18:11614–11623.
- Grossi V, Raphael D. 2003. Long-chain (C19–C29) 1-chloro-*n*-alkanes in leaf waxes of halophytes of the Chenopodiaceae. *Phytochemistry.* 63:693–698.
- Huang X, Wang C, Zhang J, Wiesenberger GLB, Zhang Z, Xie S. 2011. Comparison of free lipid compositions between roots and leaves of plants in the Dajiuhu Peatland, central China. *Biochem J.* 45:365–373.
- Joo T, Sowndhararajan K, Hong S, Lee J, Park SY, Kim S, Jhoo JW. 2014. Inhibition of nitric oxide production in LPS-stimulated RAW 264.7 cells by stem bark of *Ulmus pumila* L. *Saudi J Biol Sci.* 21:427–435.
- Kawai Y, Kumagai H, Kurihara H, Yamazaki K, Sawano R, Inoue N. 2006. β -Glucosidase inhibitory activities of phenylpropanoid glycosides, vanicoside A and B from *Polygonum sachalinense* rhizome. *Fitoterapia.* 77:456–459.
- Kazantzoglou G, Magiatis P, Kalpoutzakis E, Skaltsounis AL. 2009. Polygonophenone, the first MEM-substituted natural product, from *Polygonum maritimum*. *J Nat Prod.* 72:187–189.
- Kilinc M, Karaer F. 1995. The vegetation of Sinop Peninsula. *Turk J Bot.* 19:107–124.
- Kim KW, Ha KT, Park CS, Jin UH, Chang HW, Lee IS, Kim CH. 2007. *Polygonum cuspidatum*, compared with baicalin and berberine, inhibits inducible nitric oxide synthase and cyclooxygenase-2 gene expressions in RAW 264.7 macrophages. *Vascul Pharmacol.* 47:99–107.
- Krentz AJ, Bailey CJ. 2005. Oral antidiabetic agents: current role in type 2 diabetes mellitus. *Drugs.* 65:385–411.
- Ksouri R, Megdiche W, Falleh H, Trabelsi N, Boulaaba M, Smaoui A, Abdelly C. 2008. Influence of biological, environmental and technical factors on phenolic content and antioxidant activities of Tunisian halophytes. *C R Biol.* 331:865–873.
- Kubes P. 2000. Inducible nitric oxide synthase: a little bit of good in all of us. *Gut* 47:6–9.
- Kwon YI, Apostolidis E, Shetty K. 2007. *In vitro* studies of eggplant (*Solanum melongena*) phenolics as inhibitors of key enzymes relevant for type 2 diabetes and hypertension. *Bioresour Technol.* 99:2981–2988.
- Lean Teik NG, Chun-Han S, Min-Nan L. inventors; Kang Jian Biotech Corp., Ltd, assignee. 2013. α -Glucosidase inhibitor. US Patent 20130165679.
- Lei X, Chen J, Ren J, Li Y, Zhai J, Mu W, Zhang L, Zheng W, Tian G, Shang H. 2015. Liver damage associated with *Polygonum multiflorum* Thunb: a systematic review of case reports and case series. *Ev Based Complement Alternat Med.* 2015:9.
- Loizou S, Lekakis I, Chrousos GP, Moutsatsou P. 2010. β -Sitosterol exhibits anti-inflammatory activity in human aortic endothelial cells. *Mol Nutr Food Res.* 54:551–558.
- Maritim AC, Sanders RA, Watkins JB. 2003. Diabetes, oxidative stress, and antioxidants: a review. *J Biochem Mol Toxicol.* 17:24–38.
- Martich GD, Boujoukos AJ, Suffredini AF. 1993. Response of man to endotoxin. *Immunobiology.* 187:403–416.
- Mathers CD, Loncar D. 2006. Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Med.* 3:e442.
- Megías C, Pastor-Cavada E, Torres-Fuentes C, Girón-Calle J, Alaiz M, Juan R, Pastor J, Vioque J. 2009. Chelating, antioxidant and antiproliferative activity of *Vicia sativa* polyphenol extracts. *Eur Food Res Technol.* 230:353–359.
- Miao H, Chen L, Hao L, Zhang X, Chen Y, Ruan Z, Liang H. 2015. Stearic acid induces proinflammatory cytokine production partly through activation of lactate-HIF1 α pathway in chondrocytes. *Sci Rep.* 5:13092.
- Miranda KM, Espey MG, Wink DA. 2001. A rapid, simple spectrophotometric method for simultaneous detection of nitrate and nitrite. *Nitric Oxide.* 5:62–71.
- Miyazawa M, Tamura N. 2007. Components of the essential oil from sprouts of *Polygonum hydropiper* L. ('Benitade'). *Flavour Frag J.* 22:188–190.
- Mosmann T. 1983. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods.* 65:55–63.
- Mozaffarian D, Kamineni A, Carnethon M, Djoussé L, Mukamal KJ, Siscovick D. 2009. Lifestyle risk factors and new-onset diabetes mellitus in older adults: the cardiovascular health study. *Arch Intern Med.* 169:798–807.
- Müller C, Riederer M. 2005. Plant surface properties in chemical ecology. *J Chem Ecol.* 1:2621–2651.
- Muriach M, Flores-Bellver M, Romero FJ, Barcia JM. 2014. Diabetes and the brain: oxidative stress, inflammation, and autophagy. *Oxid Med Cell Longev.* 102:158.
- Nishishiro M, Arikawa S, Wakabayashi H, Hashimoto K, Saton K, Yokoyama K, Unten S, Kakuta H, Kurihara T, Motohashi N, et al. 2005. Inhibition of LPS-stimulated NO production in mouse macrophage-like cells by azule-nequinones. *Anticancer Res.* 25:4157–4163.
- Oh YT, Lee JY, Lee J, Lee JH, Kim JE, Ha J. Kang 2010. Oleamide suppresses lipopolysaccharide-induced expression of iNOS and COX-2 through inhibition of NF-kappaB activation in BV2 murine microglial cells. *Neurosci Lett.* 3:148–153.
- Oki T, Matsui T, Osajima Y. 1999. Inhibitory effect of α -glucosidase inhibitors varies according to its origin. *J Agric Food Chem.* 47:550–553.
- Ortiz-Andrade RR, García-Jiménez S, Castillo-España P, Ramírez-Avila G, Villalobos-Molina R, Estrada-Soto S. 2007. α -Glucosidase inhibitory activity of the methanolic extract from *Tournefortia hartwegiana*: an anti-hyperglycemic agent. *J Ethnopharmacol.* 109:48–53.
- Pan Y, Zhang X, Wang H, Liang Y, Zhu J, Li H, Zhang Z, Wu Q. 2007. Antioxidant potential of ethanolic extract of *Polygonum cuspidatum* and application in peanut oil. *Food Chem.* 105:1518–1524.
- Pereira DM, Vinholes J, Pinho PG, Valentão P, Mouga T, Teixeira N, Andrade PB. 2012. A gas chromatography-mass spectrometry multi-target method for the simultaneous analysis of three classes of metabolites in marine organisms. *Talanta.* 100:391–400.
- Ren J, Chung SH. 2007. Anti-inflammatory effect of α -linolenic acid and its mode of action through the inhibition of nitric oxide production and inducible nitric oxide synthase gene expression via NF-kappaB and mitogen-activated protein kinase pathways. *J Agric Food Chem.* 55:5073–5080.
- Richard D, Kef K, Barbe U, Bausero P, Visioli F. 2008. Polyunsaturated fatty acids as antioxidants. *Pharmacol Res.* 57:451–455.
- Rodrigues MJ, Gangadhar KN, Vizetto-Duarte C, Wubshet SG, Nyberg NT, Barreira L, Varela J, Custódio L. 2014. Maritime Halophyte species from southern Portugal as sources of bioactive molecules. *Mar Drugs.* 12:2228–2244.
- Rodrigues MJ, Soszynski A, Martins A, Rauter AP, Neng NR, Nogueira JMF, Varela J, Barreira L, Custódio L. 2015. Unravelling the antioxidant potential and the phenolic composition of different anatomical organs of the marine halophyte *Limonium algarvense*. *Ind Crops Prod.* 77:315–322.

- Ruhe RC, McDonald RB. 2001. Use of antioxidant nutrients in the prevention and treatment of type 2 diabetes. *J Am Coll Nutr.* 20:363S–369S.
- Sabu MC, Kuttan R. 2002. Anti-diabetic activity of medicinal plants and its relationship with their antioxidant property. *J Ethnopharmacol.* 81:155–160.
- Saïdana D, Mahjoub S, Boussaada O, Chriaa J, Mahjoub MA, Chéraif I, Daami M, Mighri Z, Helal AN. 2008. Antibacterial and antifungal activities of the essential oils of two saltcedar species from Tunisia. *J Am Oil Chem Soc.* 85:817–826.
- Santos CCMP, Salvadori MS, Mota VG, Costa LM, Almeida AAC, Oliveira GAL, Costa JP, Sousa DP, Freitas RM, Almeida RN. 2013. Antinociceptive and antioxidant activities of phytol *in vivo* and *in vitro* models. *Neurosci J.* 2013:9.
- Shai LJ, Magano SR, Lebelo SL, Mogale AM. 2011. Inhibitory effects of five medicinal plants on rat alpha-glucosidase: comparison with their effects on yeast alpha-glucosidase. *J Med Plants Res.* 5:2863–2867.
- Shinde J, Taldone T, Barletta M, Kunaparaju N, Hu B, Kumar S, Placido J, Zito SW. 2008. alpha-Glucosidase inhibitory activity of *Syzygium cumini* (Linn.) Skeels seed kernel *in vitro* and in Goto-Kakizaki (GK) rats. *Carbohydr Res.* 343:1278–1281.
- Soumyanath A. 2005. Traditional Medicines for Modern Times: Antidiabetic plants. New York: CRC Press, Taylor and Francis Group.
- Su CH, Hsu CH, Ng LT. 2013. Inhibitory potential of fatty acids on key enzymes related to type 2 diabetes. *Biofactors.* 39:415–421.
- Takasaki M, Konoshima T, Kuroki S, Tokuda H, Nishino H. 2001. Cancer chemopreventive activity of phenylpropanoid esters of sucrose, vanicoside B and lapathoside A, from *Polygonum lapathifolium*. *Cancer Lett.* 173:133–138.
- Trabelsi N, Falleh H, Jallali I, Daly AB, Hajlaoui H. 2012. Variation of phenolic composition and biological activities in *Limoniastrum monopetalum* L. organs. *Acta Physiol Plant.* 34:87–96.
- Valdearcos M, Esquinas E, Meana C, Peña L, Gil-de-Gómez L, Balsinde J, Balboa MA. 2012. Lipin-2 reduces proinflammatory signaling induced by saturated fatty acids in macrophages. *J Biol Chem.* 287:10894–10904.
- Vassiliou EK, Gonzalez A, Garcia C, Tadros JH, Chakraborty G, Toney JH. 2009. Oleic acid and peanut oil high in oleic acid reverse the inhibitory effect of insulin production of the inflammatory cytokine TNF-alpha both *in vitro* and *in vivo* systems. *Lipids Health Dis.* 8:25.
- Veerasamy R, Min LS, Pauine R, Sivadasan S, Varghese C, Rajak H, Marimuthu K. 2014. Effect of aqueous extract of *Polygonum minus* leaf on the immunity and survival of African catfish (*Clarias gariepinus*). *J Coast Life Med.* 2:209–213.
- Vikram A, Tripathi DN, Kumar A, Singh S. 2014. Oxidative stress and inflammation in diabetic complications. *Int J Endocrinol.* 2014:2.
- Wink DA, Hines HB, Cheng RY, Switzer CH, Flores-Santana W, Vitek MP, Ridnour LA, Colton CA. 2011. Nitric oxide and redox mechanisms in the immune response. *J Leukoc Biol.* 89:873–891.
- World Health Organization (WHO). 1999. Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: diagnosis and classification of diabetes mellitus. Geneva (WHO/NCD/NCS/99.2).
- Xiao Z, Storms R, Tsang A. 2006. A quantitative starch-iodine method for measuring alpha-amylase and glucoamylase activities. *Anal Biochem.* 351:146–148.
- Yarchoan M, Arnold SE. 2014. Repurposing diabetes drugs for brain insulin resistance in Alzheimer disease. *Diabetes.* 63:2253–2261.