



## Complete mitochondrial genomes of three species of fresh flies of forensic entomology interest from the genus *Sarcophaga* (Sarcophagidae) from Portugal and Brazil

Carmo Anderson Oliveira, Susanne Faccin, Patrícia J. Thyssen, Deodália Dias, Evanguedes Kalapothakis & Maria Teresa Rebelo

To cite this article: Carmo Anderson Oliveira, Susanne Faccin, Patrícia J. Thyssen, Deodália Dias, Evanguedes Kalapothakis & Maria Teresa Rebelo (2019) Complete mitochondrial genomes of three species of fresh flies of forensic entomology interest from the genus *Sarcophaga* (Sarcophagidae) from Portugal and Brazil, Mitochondrial DNA Part B, 4:1, 237-239, DOI: [10.1080/23802359.2018.1545539](https://doi.org/10.1080/23802359.2018.1545539)

To link to this article: <https://doi.org/10.1080/23802359.2018.1545539>



© 2018 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 26 Dec 2018.



Submit your article to this journal [↗](#)



Article views: 12



View Crossmark data [↗](#)

MITOGENOME ANNOUNCEMENT



## Complete mitochondrial genomes of three species of fresh flies of forensic entomology interest from the genus *Sarcophaga* (Sarcophagidae) from Portugal and Brazil

Carmo Anderson Oliveira<sup>a,b</sup> , Susanne Faccin<sup>a</sup> , Patrícia J. Thyssen<sup>c</sup> , Deodália Dias<sup>b,d</sup> , Evanguedes Kalapothakis<sup>a</sup>  and Maria Teresa Rebelo<sup>b,d</sup> 

<sup>a</sup>Laboratório de Biotecnologia e Marcadores Moleculares, Universidade Federal de Minas Gerais, UFMG, Brazil; <sup>b</sup>CESAM – Centro de Estudos do Ambiente e do Mar Faculdade de Ciências da Universidade de Lisboa, Portugal; <sup>c</sup>Departamento de Biologia Animal, Universidade Estadual de Campinas, UNICAMP, Brazil; <sup>d</sup>Departamento de Biologia Animal Faculdade de Ciências da Universidade de Lisboa, Portugal

### ABSTRACT

The Sarcophagidae family of fresh flies bears strong importance in the context of forensic entomology due to their application in the estimation of the Post Mortem Interval (PMI). *Sarcophaga* is the major genus in the Sarcophagidae family and includes cosmopolitan species, which are distributed worldwide. In this communication, we present the analysis of the complete mitochondrial genome (mtDNA) of two species from Portugal – *S. melanura* and *S. dux* – and one from Brazil – *S. ruficornis*. The mtDNA of these species range from 14,882 bp to 15,190 bp and have 22 tRNA genes, 13 protein-coding genes (PCG), and two rRNAs distributed along both strands. Our data include the first record of complete *Sarcophaga* mtDNA sequences from species collected in Portugal and in Brazil. These genomes represent an advance in the understanding about this group, expand the database, and can be used for the development of new markers for species identification.

### ARTICLE HISTORY

Received 15 October 2018  
Accepted 2 November 2018

### KEYWORDS

Forensic entomology;  
*Sarcophaga*; NGS; complete  
mtDNA; flesh flies

Sarcophagidae is one of the most important families in forensic entomology due to its widespread distribution and diversity of species (Pape 1996). The genus *Sarcophaga* is considered the most diverse, with 133 subgenera and 790 species described (Pape 1996). Species belonging to this genus are frequently used and represent an important source of information in forensic investigations (Moura et al. 1997; Carvalho et al. 2000; Oliveira et al., 2011; Rolo et al., 2013). The correct identification of species from the *Sarcophaga* genus is usually made using the male genitalia. However, in contrast with other genus, the females of some *Sarcophaga* species can be identified by their external morphology (Zhang et al. 2014). Nevertheless, morphological identification still presents its limitations, which hinder the identification of larval stages and dandified specimens. We show novel complete mtDNA sequences of flesh fly species of forensic importance. Our data expand the knowledge on the molecular database for the identification of these species.

In this work, we collected *Sarcophaga* specimens from two sites in Portugal and one in Brazil. In Portugal, *S. dux* (38°35'02" N 9°10'30" W), and *S. melanura* (38°42'52" N 9°11'45" W). *Sarcophaga ruficornis* was collected in Brazil (22°51'12" S 46°19'18" W). Each specimen was identified following the taxonomic key (Buenaventura and Pape 2015) and preserved in 70% ethanol. The mtDNA was extracted from the thorax, legs, and wings of the flies (Françoso et al.


2016), and the head and abdomen of *S. ruficornis* (Voucher 14) were deposited in the database of UNICAMP, whereas *S. dux* (Voucher 2) and *S. melanura* (Voucher GE) were registered in the database of the Universidade de Lisboa. The mtDNA library was generated using the Nextera XT kit according to the manufacturer's instructions and sequenced using a paired-end strategy 2 × 250 on the MiSeq platform (Illumina, CA). The genomes were constructed by mapping the reads against the *Sarcophagidae* mitogenomes available on the NCBI database, followed by *de novo* assembly of the mapped reads using the CLC Genomics Workbench. The mitogenomes were annotated using the MITOS WebServer (Bernt et al. 2013) and manually verified using the NCBI database.

The complete mtDNA of *S. ruficornis* (MH879755) was 14,940 bp long and presented 24.3% of CG. *S. dux* (MH879759) mtDNA was 14,882 bp long and showed 23.8% of CG. *S. melanura* (MH879758) presented the longest mtDNA, with 15,190 bp and 24.4% of CG. The annotation of these genomes revealed 13 protein-coding genes (PCGs), two rRNA genes, 22 tRNA genes, and a noncoding control region (D-loop) located between the 12S rRNA and tRNA<sup>Ile</sup>. The PCG sequences commonly start with an ATT, ATA, or ATG codon (12 PCGs), but COX1 starts with CAA. Five PCGs have the T-stop codon completed to TAA by posttranscriptional polyadenylation (Ojala et al. 1981). The light strand codifies eight of the 22 tRNAs, five PCGs and the two rRNAs, while the

**CONTACT** Evanguedes Kalapothakis  [kalapothakis@gmail.com](mailto:kalapothakis@gmail.com)  Av. Antônio Carlos, 6627, Pampulha, Belo Horizonte, Minas Gerais, Brazil

© 2018 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



**Figure 1.** Molecular phylogenetic inferences of *Sarcophaga* mitogenomes. The phylogenetic analyses were made using the MEGA 7 software (Kumar et al. 2016) using the maximum likelihood model with the Nearest-Neighbour-Interchange heuristic method. The D-loop region was excluded from this analysis due to its high degree of variability (Gonder et al., 2007). The tree showed a well-structured separation of the novel species. *S. melanura* (collected in Lisbon, Portugal) showed few differences in comparison with the species from NCBI collected in China (KP091687.1). *S. dux* (Aroeira, Portugal) is more related to *S. portschinskyi* (China). On the other hand, *S. ruficornis* (Brazil) clustered with the branch including *S. crassipalpis* and *N. bullata* (both from the United States). The novel mtDNA published herein are highlighted in grey. The mitogenomes for comparison were obtained from the NCBI database: *Musca domestica* (KT272857.1) – used as outgroup, *Sarcophaga impatiens* (JN859549.1), *Sarcophaga melanura* (KP091687.1), *Neobellieria bullata* (KT272859.1), *Sarcophaga crassipalpis* (KC005711.1), *Sarcophaga peregrina* (KF921296.1), *Sarcophaga formosensis* (MF688648.1), *Sarcophaga africa* (KM881633.1), *Sarcophaga portschinskyi* (KM287570.1), *Sarcophaga similis* (KM287431.1), *Sarcophaga albiceps* (KT444443.1), *Sarcophaga misera* (MF133500.1).

remaining genes are encoded on the heavy strand. The phylogenetic analysis and comparison with others *Sarcophaga* species are showed in Figure 1.

Deodália Dias <http://orcid.org/0000-0002-8771-6727>  
 Evanguedes Kalapothakis <http://orcid.org/0000-0002-8326-249X>  
 Maria Teresa Rebelo <http://orcid.org/0000-0002-2724-2195>

## Acknowledgments

We thank the Universidade de Lisboa and the Universidade Federal de Minas Gerais (UFMG) for the support during the project execution.

## Disclosure statement

The authors report no potential conflict of interest.

## Funding

This work was funded by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) (Edital Ciências Forenses no. 25/2014, Process 23038.006839 / 2014-33), International postdoc fellowship (CAPES 0092/17-7) and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq). Financial support was also provided by CESAM (UID/AMB/50017 - POCL-01-0145-FEDER-007638), and by FCT/MCTES through national funds (PIDDAC). Co-funding was provided by the FEDER, within the PT2020 Partnership Agreement and Compete 2020.

## ORCID

Carmo Anderson Oliveira <http://orcid.org/0000-0003-4646-513X>  
 Susanne Faccin <http://orcid.org/0000-0003-2145-0396>  
 Patrícia J. Thyssen <http://orcid.org/0000-0001-7343-2419>

## References

- Bernt M, Donath A, Jühling F, Externbrink F, Florentz C, Fritzsch G, Pütz J, Middendorf M, Stadler PF. 2013. MITOS: improved de novo metazoan mitochondrial genome annotation. *Mol Phylogenet Evol.* 69: 313–319.
- Buenaventura E, Pape T. 2015. Phylogeny of the Peckia-Genus group: evolution of male genitalia in the major Necrophagous guild of neotropical flesh flies (Diptera: Sarcophagidae). *Org Divers Evol.* 15: 301–331.
- Carvalho L, Thyssen PJ, Linhares A, Palhares F. 2000. A checklist of Arthropods associated with pig carrion and human corpses in southeastern Brazil. *Memorias Do Instituto Oswaldo Cruz.* 95: 135–138.
- Françoso E, Gomes F, Arias MC. 2016. A protocol for isolating insect mitochondrial genomes: a case study of NUMT in *Melipona flavolineata* (Hymenoptera: Apidae). *Mitochondrial DNA.* 27: 2401–2404.
- Gonder MK, Mortensen HM, Reed FA, Sousa A, Tishkoff SA. 2007. Whole-MtDNA genome sequence analysis of ancient African lineages. *Mol Biol Evol.* 24:757–768. doi:10.1093/molbev/msl209.
- Kumar S, Stecher G, Tamura K. 2016. MEGA7: molecular evolutionary genetics analysis version 7. 0 for bigger datasets. *Mol Evol Genet Anal.* 33: 1870–1874.
- Moura MO, Carvalho CJB, Monteiro-Filho ELA. 1997. A preliminary analysis of insects of medico-legal importance in Curitiba, State of Paraná, *Memorias. Do Instituto Oswaldo Cruz.* 92:269–274.

- Ojala D, Montoya J, Attardi G. 1981. tRNA punctuation model of RNA processing in human mitochondria. *Nature*. 290:470.
- Oliveira AR, Farinha A, Rebelo MT, Dias D. 2011. Forensic entomology: molecular identification of blowfly species (Diptera: Calliphoridae) in Portugal. *Forensic Sci Int*. 3:e439–e440.
- Pape T. 1996. Catalogue of the Sarcophagidae of the world (Insecta: Diptera). *Memoirs Entomol Int*. 8:1–558.
- Rolo EA, Oliveira AR, Dourado CG, Farinha A, Rebelo MT, Dias D. 2013. Identification of sarcosaprophagous Diptera species through DNA barcoding in wildlife forensics. *Forensic Sci Int*. 228: 160–164.
- Zhang M, Chen JL, Gao XZ, Pape T, Zhang D. 2014. First description of the female of *Sarcophaga* (*Sarcophaga*) *gracilior* (Chen, 1975) (Diptera, Sarcophagidae). *Zookeys*. 2:43–53.