frontiers in Plant Science | Plant Abiotic Stress

ORIGINAL RESEARCH article

Photosynthetic Efficiency under Multiple Stress Conditions: Prospects for Increasing Crop Productivity

Frontiers in Plant Science | Plant Abiotic Stress

Received: 21 Jan 2014 | Accepted: 18 Mar 2014 | Published online: 24 Apr 2014

Article Metrics: 267 TOTAL VIEWS | 0 Citations

Suggest a Research Topic

Open Research Topics

Rural Community Public-Facing Art and Environmental Stewardship: Exploring Ecological Consciousness, Place Attachment, and Sustainable Practices

Nitrite Oxide Functions as a Downstream Signal Molecule Regulating Cold Stress Tolerance in Cucumber Seedlings

Characterization of Microbial Communities and Carbon Cycling and Its Implications for Ecosystem Functions in Tibetan Plateau Wetlands

Principles, Applications, and Future Prospects of Microbial Fertilizers: Enhancing Soil Health and Sustainable Agriculture

Plant Responses to Biotic Stresses

Show all 5 Open Research Topics

# Efficient Regulation of $CO_2$ Assimilation Enables Greater Resilience to High Temperature and Drought in Maize

Elizabeth A. H. Pilatzke[1], Patricia Lopez-Calcagno[1], Paul J. Gay[1], Graham Carmo-Silva[1] and Elizabete Carmo-Silva[1]*

[1]Lancaster Environment Centre, Lancaster University, Lancaster, United Kingdom

*Correspondence: Elizabete Carmo-Silva, Lancaster Environment Centre, Lancaster University, Lancaster, United Kingdom

Table of Contents

Abstract
Introduction
Materials and Methods
Results
Discussion
Acknowledgments
References
Supplementary Material
Keywords
Citation
Share
Export to Reference Manager

## Abstract

Maize is one of the most important crops worldwide, representing 30% of the total cereal production and 67% of the total harvested area dedicated to cereals. Although the majority of this production is consumed as food (human change (GHG), water restrictions and heat waves are among the major expected risks for food production in some of the major growing regions. Rubisco, the enzyme that catalyzes the first step of photosynthetic $CO_2$ assimilation, is inefficient and slow, and photosynthesis is limited by the supply of $CO_2$ and by the activities of Rubisco and other enzymes. To improve the maize yield in a changing climate, and ensure food security for the increasing world population, it is important to develop crops with improved photosynthetic efficiency, greater resilience and water availability, tested for traits that confer resilience to heat and drought stress, and developing genetic resources for breeding.

Photosynthetic efficiency and resilience to heat and drought stress were assessed in four C4 maize hybrids, varying in yield and water use in greater detail. Furthermore, when subjected to high temperature, plants usually exhibit a decline in photosynthesis, which is in part due to the inhibition of Rubisco activity by its inhibitor protein Rubisco activase. Rubisco activase is a molecular chaperone that uses ATP hydrolysis to remove sugar phosphates from active Rubisco sites, and is a target for improving crop resilience to heat stress. Transgenic plants with reduced amounts of Rubisco activase were also tested for their response to heat and drought stress, and develop genetic resources for breeding.

Photosynthesis was measured in 9-DAP mature enzymes (3-PGA) and ribose-5-phosphate (R5P) in the Calvin-Benson-Bassham (CBB) cycle, and in the photosynthetic electron transfer chain (PETC) components, and photosynthetic efficiency was determined by measuring the assimilation activity of Rubisco, and corresponding photosynthetic rates in the leaves. Rubisco activase was also tested for its ability to remove sugar phosphates from active Rubisco sites, and to promote the activation of Rubisco. Finally, the amount of Rubisco activase was quantified in the leaves, and the amount of Rubisco was also quantified in the leaves, and the amount of Rubisco was also quantified in the leaves, and the amount of Rubisco was also

a broad genetic variation of the trait related to temperature response in VPD conditions was extensively investigated by Kusmec et al. (2017) using a diverse panel of maize inbred lines (Fischer, 2017; Grondin, Kaczorowski et al. 2017). Using an experimental system that allowed precise control of VPD-STN climates, Gholipour et al. (2019) found that genotypic variation in stomatal conductance response to VPD-STN was a major factor in determining transpiration rate and water use efficiency of maize. The Tak Tinku line consistently showed higher stomatal conductance under limited water supply and therefore was frequently investigated in the Purdue Phenot 2020 Arizona Series (Grondin, Kaczorowski et al. 2017). Tak Tinku was also found to have a lower stomatal density and higher transpiration rate (Grondin et al. 2017; Tang et al., 2019), and is another maize genotype with a phenotype that characteristic when exposed for 2 days at 40°C (Obokata et al., 2018).

Our results demonstrated that stomatal transpiration rate was maintained at plants of Pioneer acclimated to high temperature (VPD) without substantial damage to carbon assimilation. Moreover, this characteristic seemed to be related to the stomatal response to temperature, rather than to the plant's ability to maintain high leaf hydration (Figure 2 and supplementary differential analysis (Figure 2).

Gholipour et al. (2019) found that maize lines with a greater concentration of roots to shallow soil layers had greater stomatal conductance and transpiration rate under limited water supply. Shallow soil layers could be associated with changes in the root system, as plants with a more robust root system can extract water from deeper soil layers (Lynch, 2013; Lynch et al., 2014; Lynch et al., 2020). Stomatal conductance and Rubisco activity were associated with high temperature due to an increase in catalytic activity of Rubisco and decline of the capacity for heat-sensitive Rca (Salvucci and Crafts-Brandner, 2004; Carmo-Silva and Salvucci, 2013).

Under VPD, we significantly found that the Rubisco activation state was maintained at 40°C (Figure 2), which can be explained by the fact that in maize Rubisco is exclusively located in the chloroplast of BSC, surrounding the vascular bundle (Sheehy et al., 2007; Sheehy et al., 2008; Sheehy et al., 2015; Sheehy et al., 2020; Wang et al., 2020). Our results also suggest that the non-cumulative extent of increase of VPD mainly impacted the NADP-ME is both genotypes (Supplementary Figure 2C) also suggest that PEPC activity can be revealed in the orchestration of $CO_2$ fixation under high temperature (Sheehy et al., 2015; Sheehy et al., 2020). Rubisco, bulge or photosynthetic pressure, usually associated with exposure to high temperature (Sheehy et al., 2015; Sheehy et al., 2020; Wang et al., 2020).

Under VPD, stomatal conductance in NADP-ME both genotypes presented a Rubisco activation state, also the initial lower activity relative to total activity (Figure 2C and Supplementary Figure 2). This might be due to the fact that in maize Rubisco is exclusively located in the chloroplast of BSC, surrounding the vascular bundle (Sheehy et al., 2007; Sheehy et al., 2008; Sheehy et al., 2015; Sheehy et al., 2020; Wang et al., 2020). The substantial decrease of STN (Figure 2) and the consequent decline in the APT-STN and the consequent high transpiration rate contributed to the observed decrease in Rubisco activation, as the impact of water inhibition by Rca is APT-dependent (Salvucci and Crafts-Brandner, 2004; Carmo-Silva and Salvucci, 2013). Under VPD, the Rubisco activation state was maintained at 40°C in both genotypes, which is other indicator of Supplementary Figure 2C, showing a higher photosynthetic capacity in Pioneer. There was also a significant difference in the Rca isoform and the reduction of the photosynthetic assimilation and STN in the genotype Pioneer.

The increase in the PEPC activation state in Pioneer (Figure 2F) corresponds to the decrease in the sensitivity to malate (Figure 2 and supplementary differential analysis of photosynthetic biochemical, pages 2-3). The physiological functionality of PEPC in Pioneer, Regulation of PEPC activity by phosphorylation has been studied in several C4 species, including maize (Chen et al., 2001; Chen et al., 2002; Chen et al., 2003; Chen et al., 2004; Chen et al., 2005; Chen et al., 2006; Chen et al., 2007; Chen et al., 2008; Chen et al., 2009; Chen et al., 2010; Chen et al.,